



# 15-0-0

## Turf King Fertilizer with 0.58% Prodiamine

**For Pre-Emergence Control Of Grass And Broadleaf Weeds In Lawns And Ornamental Turf (Excluding Golf Course Putting Greens)**

**Landscape Ornamentals, Established Perennials And Wildflower Plantings**

**Stops Many Weed Grasses Before They Grow**

**Prevents Crabgrass, Annual Bluegrass, Foxtail**

**ACTIVE INGREDIENT:**  
 Prodiamine: (CAS # 29091-21-2) ..... 0.58%

**INERT INGREDIENTS:** ..... 99.42%

**TOTAL:** ..... 100.00%

### KEEP OUT OF REACH OF CHILDREN CAUTION

See back panel for First Aid, Precautionary Statements, Directions for Use and Storage and Disposal.

### PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS CAUTION

Causes moderate eye injury (irritation). Harmful if inhaled or absorbed through the skin. Avoid contact with skin, eyes, or clothing. Avoid breathing dust. Wash thoroughly with soap and water after handling. Prolonged or repeated skin contact with product may cause allergic reactions in some individuals.

#### FIRST AID

|                                |  |
|--------------------------------|--|
| <b>IF IN EYES:</b>             | <ul style="list-style-type: none"> <li>Hold eye open and rinse slowly and gently with water for 15-20 minutes.</li> <li>Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li> <li>Call a poison control center or doctor for treatment advice.</li> </ul>  |
| <b>IF ON SKIN OR CLOTHING:</b> | <ul style="list-style-type: none"> <li>Take off contaminated clothing.</li> <li>Rinse skin immediately with plenty of water for 15-20 minutes.</li> <li>Call a poison control center or doctor for treatment advice.</li> </ul>  |
| <b>IF SWALLOWED:</b>           | <ul style="list-style-type: none"> <li>Call a poison control center or physician immediately for treatment advice.</li> <li>Have person sip a glass of water if able to swallow.</li> <li>Do not induce vomiting unless told to do so by the poison control center or physician.</li> <li>Do not give anything by mouth to an unconscious person.</li> </ul> |
| <b>IF INHALED:</b>             | <ul style="list-style-type: none"> <li>Move person to fresh air.</li> <li>If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible.</li> <li>Call a poison control center or doctor for further treatment advice.</li> </ul>   |

#### HOT LINE NUMBER

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact SafetyCall® International (866) 897-8050 for emergency medical treatment information.

#### ENVIRONMENTAL HAZARDS

This product has low solubility in water. At the limits of solubility, this product is not toxic to fish. However, at concentrations above the level of water solubility, it may be toxic to fish. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Drift and runoff from treated areas may be hazardous to aquatic organisms in adjacent sites. Do not contaminate water when disposing of equipment washwaters.

Distributed and Guaranteed by Andre & Son  
 17150 State Route 706 • Montrose, PA 18801 • 570-278-1131

#### GUARANTEED ANALYSIS

Total Nitrogen (N) ..... 15.00%  
 10.5% Urea Nitrogen  
 4.5% CSRUN

Sulfur (S) ..... 0.40%

**Nutrients derived from:**  
 XCU, Urea.

Information concerning the raw materials composing this product can be obtained by writing to the Guarantor listed on this label, and referring to the batch number found on this container.

Information regarding the contents and levels of metals in this product is available on the internet at <http://www.aapfco.org/metals.htm>

#### WEED RESISTANCE MANAGEMENT

For resistance management, Turf King Fertilizer with 0.58 Prodiamine is a Group 3 herbicide. Any weed population may contain or develop plants naturally resistant to Turf King Fertilizer with 0.58 Prodiamine and other Group 3 herbicides. The resistant biotypes may dominate the weed population if these herbicides are used repeatedly in the same field. Appropriate resistance-management strategies should be followed.

To delay herbicide resistance, take one or more of the following steps:

- Rotate the use of Turf King Fertilizer with 0.58 Prodiamine or other Group 3 herbicides within a growing season sequence or among growing seasons with different herbicide groups that control the same weeds in a field.
- Use tank mixtures with herbicides from a different group if such use is permitted; where information on resistance in target weed species is available, use the less resistance-prone partner at a rate that will control the target weed(s) equally as well as the more resistance prone partner. Consult your local extension service or certified crop advisor if you are unsure as to which active ingredient is currently less prone to resistance.
- Adopt an integrated weed-management program for herbicide use that includes scouting and uses historical information related to herbicide used and crop rotation, and that considers tillage (or other mechanical control methods), cultural (e.g. higher crop seeding rates; precision fertilizer application method and timing to favor the crop and not the weeds), biological (weed-competitive crops or varieties) and other management practices.
- Scout after herbicide application to monitor weed populations for early signs of resistance development. Indicators of possible herbicide resistance include: (1) failure to control a weed species normally controlled by the herbicide at the dose applied, especially if control is achieved on adjacent weeds; (2) a spreading patch of non-controlled plants of a particular weed species; (3) surviving plants mixed with controlled individuals of the same species. If resistance is suspected, prevent weed seed production in the affected area by an alternative herbicide from a different group or by a mechanical method such as hoeing or tillage. Prevent movement of resistant weed seeds to other fields by cleaning harvesting and tillage equipment when moving between fields, and planting clean seed.
- If a weed pest population continues to progress after treatment with this product, discontinue use of this product, and switch to another management strategy or herbicide with a different mode of action, if available.
- Contact your local extension specialist or certified crop advisors for additional pesticide resistance-management and/or integrated weed-management recommendations for specific crops and weed biotypes.
- Report any incidence of non-performance of this product against a particular weed species to Control Solutions, Inc., your local retailer or your local extension specialist. If resistance is suspected, treat weed escapes with a herbicide having a different mechanism of action and/or use non-chemical means to remove escapes, as practical, with the goal of preventing further seed production.

#### BEST MANAGEMENT PRACTICES

Proactively implementing diversified weed control strategies to minimize selection for weed populations resistant to one or more herbicides is recommended. A diversified weed management program may include the use of multiple herbicides with different modes of action and overlapping weed spectrum with or without tillage operations and/or other cultural practices. Research has demonstrated that using the labeled rate and directions for use is important to delay the selection for resistant weeds. Scouting after herbicide application is important because it can facilitate the early identification of weed shifts and/or weed resistance and thus provide direction on future weed management practices. One of the best ways to contain resistant weed populations is to implement measures to avoid allowing weeds to reproduce by seed or to proliferate vegetatively. Cleaning equipment between sites and avoiding movement of plant material between sites will greatly aid in reducing the spread of resistant weed seed.

#### DIRECTION FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Not for use on plants being grown for sale or other commercial use, for commercial seed production, or for research purposes. For use on plants intended for aesthetic purposes or climatic modification and being grown in ornamental gardens or parks, or on golf courses or lawns and grounds.

Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes.

## NET WT 50 lb (22.7 kg)

EPA Reg. No. 53883-171-47956 • EPA Est. No. 47956-PA-001

PRODIAMINE GROUP 3 HERBICIDE

**FAILURE TO FOLLOW THE DIRECTIONS FOR USE AND PRECAUTIONS ON THIS LABEL MAY RESULT IN POOR WEED CONTROL, CROP INJURY, OR ILLEGAL RESIDUES.**

This product is a pre-emergence herbicide that provides residual control of many grass and broadleaf weeds in established lawns and ornamental turf (excluding golf course putting greens), landscape ornamentals and established perennials and wildflower plantings. It controls susceptible weeds by inhibiting weed seeds, germination and root development. When used according to directions, it will control; barnyardgrass; bluegrass; annual (Poa annua); carpetweed; chickweed, common; chickweed, mouseear (from seed); crabgrass (large, smooth); crowfootgrass; cupgrass, woolly; foxtails, annual; goosegrass<sup>1</sup>; henbit; itchgrass; johnsongrass (from seed); junglerice; knotweed; kochia; lambsquarter, common; lovegrass; panicum (Texas, fall, browntop); pigweed; purslane, common; pusley, Florida; rescuegrass<sup>2</sup>, shepherd's purse<sup>2</sup>, signalgrass, broadleaf; speedwell, Persian; sprangletop; spurge, prostrate; witchgrass; woodsorrel, yellow (from seed).

<sup>1</sup>In many areas a single application of 2.5 to 6 pounds/1,000 square feet (112 to 259 pounds/acre) will control goosegrass. However, under heavy goosegrass pressure and/or an extended growing season, most effective weed control may be obtained with an initial application of 2.5 to 3.9 pounds/1,000 square feet (112 to 173 pounds/acre), followed after 60 to 90 days by a second application at doses that would not exceed the maximum annual rate (see "Minimum and Maximum Application Rate by Turfgrass Species" table in the Rates of Application section of this label).


<sup>2</sup>For this weed, apply in late summer, fall or winter prior to germination.

<sup>3</sup>Suppression only. Sequential applications may be made so long as the total amount of product applied does not exceed the maximum annual application rates specified for each turf species. All applications must be made prior to germination of the weed seeds.

#### WHEN TO APPLY TO ESTABLISHED TURF:

This product must be applied prior to germination of weed seeds, as it will not control established weeds.

#### THIS MAP SHOWS APPROXIMATE CRABGRASS GERMINATION DATES



#### APPLICATION DIRECTIONS:

Apply this product uniformly, using a suitable spreader that has been properly calibrated. Most effective weed control will be obtained when this product is activated by at least 0.5 inch of rainfall or irrigation, or with shallow (1-2 inches) incorporation, prior to weed seed germination and within 14 days application.

#### USE PRECAUTIONS:

The following precautions apply to the use of this product in turf grasses and lawns. Application of this product may thin emerged annual bluegrass and newly overseeded grasses.

- Application to turf stressed by drought, low fertility, or pest damage may result in turf injury.
- Disturbing the herbicide barrier with cultural practices (such as core aeration) may result in reduced weed control.

#### USE RESTRICTIONS:

- Do not apply to overseeded turf within 60 days after seeding or until after the second mowing, whichever is longer. Injury to desirable seedlings is likely if this product is applied before seedling secondary roots are in the second inch of soil (not thatch plus soil).
- Do not apply to newly set sod until the following year.
- Do not apply to putting greens, or areas where dichondra, colonial bentgrass, velvet bentgrass or annual bluegrass (*Poa annua*) are desirable species.
- Do not apply through any type of irrigation system.
- Do not apply aerially.
- Do not graze or feed livestock forage cut from treated areas.

#### RATES OF APPLICATION:

This product is recommended for use on the turf grass species listed in the following table. The maximum amount of this product that may be applied per year is given for each turf grass species. Do not exceed the maximum yearly rate listed:

| MINIMUM AND MAXIMUM APPLICATION RATES (by Turf Grass Species)  |  |   |
|--|--|---|
| Turf Species   | Minimum Rate                             | Maximum Annual Rate <sup>1</sup>                          |
| Bahiagrass, Bermudagrass <sup>2</sup><br>Centipedegrass, Zoysiagrass,<br>Seashore Paspalum,<br>St. Augustinegrass Tall Fescue<br>(including turf-Type) | 4 lbs/1,000 sq. ft.<br>(175 lbs/acre)    | 6 lbs/1,000 sq. ft. per year<br>(259 lbs/acre per year)   |
| Buffalograss<br>Kentucky Bluegrass<br>Perennial Ryegrass   | 2 lbs/1,000 sq. ft.<br>(87.5 lbs/acre)   | 4 lbs/1,000 sq. ft. per year<br>(173 lbs/acre per year)   |
| Creeping Red Fescue  | 1.5 lbs/1,000 sq. ft.<br>(65.3 lbs/acre) | 2.9 lbs/1,000 sq. ft. per year<br>(130 lbs/acre per year) |
| Creeping Bentgrass   | 1.5 lbs/1,000 sq. ft.<br>(65.3 lbs/acre) | 2.5 lbs/1,000 sq. ft. per year<br>(112 lbs/acre per year) |

